

NUTRIBION Trial Report

Nanda Feeds Pvt Ltd, Bangalore

Scientific field trial was conducted at Nanda Feeds Bangalore. NUTRIBION was used in one feed and another feed was regular control feed without NUTRIBION. NUTRIBION was added @ 200g in Pre Starter ration, 300g in Starter ration and 400g in Finisher ration.

Trial was conducted during June 2016 – July 2016. The details are in the tables below:

Batch No: 4B	Induction Date: 04 June 2016	
Particulars	Control Feed	With NUTRIBION
Day old broiler chicks	7627	6734
No. of days reared	42	42
Total number of birds culled	7327	6429
Total number of kgs	16557.7	14713.2
Average body weight	2.26	2.29
Mortality	300	305
Mortality %	3.93%	3.51%
FCR	1.72	1.70

Conclusion:

It has be observed that the better FCR and Better body weight was achieved in the NUTRIBION group

NUTRIBION Trial Report

Nanda Feeds Pvt Ltd, Bangalore

Scientific field trial was conducted at Nanda Feeds Bangalore. NUTRIBION was used in one feed and another feed was regular control feed without NUTRIBION. NUTRIBION was added @ 200g in Pre Starter ration, 300g in Starter ration and 400g in Finisher ration.

Trials was conducted during May 2016 – June 2016. The details are in the tables below:

Particulars	Induction Date: 20 May 2016	
	Control Feed	With NUTRIBION
No. of days reared	41	41
Total number of birds culled	5705	5492
Total number of kgs	12131.9	11982.5
Average body weight	2.13	2.18
Mortality	227	227
Mortality %	3.83%	3.81%
FCR	1.68	1.64

Conclusion:

We see a good FCR and a higher average body weight in the NUTRIBION group.